

FROM PCM TO KDV AND BACK

Jonathan M. Evans

*Joseph Henry Laboratories, Princeton University, Princeton NJ 08544, USA**DAMTP, University of Cambridge, Silver Street, Cambridge CB3 9EW, UK*

Contribution to the NATO-ASI meeting

Integrable Hierarchies and Modern Physical Theories

University of Illinois at Chicago, July 2000

This is a summary of progress made [1-4] in understanding the occurrence and properties of local, conserved, commuting charges in non-linear sigma-models, including principal chiral models (PCMs) and WZW models. In each PCM or WZW model with target manifold a compact Lie group G , there are infinitely many commuting charges whose currents have the form $\mathcal{K}_m = k_{a_1 a_2 \dots a_m} j^{a_1} j^{a_2} \dots j^{a_m}$. The underlying fields j^a take values in the Lie algebra $\mathfrak{g}$ (indices a refer to some basis) and each tensor $k_{a_1 a_2 \dots a_m}$ is totally symmetric and G -invariant. The spins of the corresponding conserved charges $s = m - 1$ run over the exponents of the algebra modulo its Coxeter number h . Such patterns of spins are familiar from the context of affine Toda field theory [5] and these similarities offer a natural explanation of some otherwise mysterious common properties of PCM and affine Toda S-matrices [1].

Initial investigations [1] focussed on PCMs based on classical groups G and established that currents and symmetric tensors with the required properties could be defined by a formula $\mathcal{K}_m = \det(1 - \mu j^a t_a)^{s/h} |_{\mu^{s+1}}$, where the generators t_a belong to the defining representation of $\mathfrak{g}$. Commutation of the resulting charges depends upon some intricate algebraic identities satisfied by the k -tensors which we shall refer to as the *commutation conditions*. These findings were subsequently extended to incorporate the effects of WZ terms and supersymmetry [2] and analogous results were also established for sigma-models whose target manifolds are symmetric spaces G/H [3], but again only for G and H classical groups. The general case, including exceptional groups, has been treated quite recently [4] by making use of a direct link between the k -tensors and the Drinfeld-Sokolov modified KdV (DS/mKdV) hierarchies [6].

The first step in establishing this link is to show that all relevant attributes of the k -tensors can be understood as properties of the *restricted* tensors $k_{i_1 i_2 \dots i_m}$ defined just on the Cartan subalgebra (CSA) of $\mathfrak{g}$ (indices i refer to the CSA). It is well-known that any symmetric invariant tensor is completely determined by its restriction and that the remnant of G which fixes the CSA is the Weyl group; moreover, any Weyl-invariant tensor on the CSA can be extended uniquely to a G -invariant tensor on $\mathfrak{g}$. It is less obvious that the key commutation conditions of specific interest to us can be simply expressed in terms of restricted tensors alone, but this is in fact the case [4].

Recall next that the DS/mKdV hierarchy associated with a Lie algebra $\mathfrak{g}$ [6] involves currents of the form $\mathcal{H}_m = h_{i_1 i_2 \dots i_m} u^{i_1} u^{i_2} \dots u^{i_m} + (\text{derivative terms})$, where the fields u^i live in the CSA of $\mathfrak{g}$. The explicit term in $\mathcal{H}_m$ contains a certain Weyl-invariant tensor $h_{i_1 i_2 \dots i_m}$, while the additional, implicit terms involve derivatives of the fields. Drinfeld and Sokolov proved that the charges corresponding to these currents always commute. It is possible to deduce from this that the h -tensors must satisfy precisely the same *commutation conditions* already mentioned, notwithstanding the complications arising from the extra derivative terms.

We can conclude, therefore, that after extension from the CSA to all of $\mathfrak{g}$, the h -tensors of the DS hierarchies provide new definitions of k -tensors with exactly the properties we require for commuting charges, for any compact Lie algebra, whether classical or exceptional. Careful examination [4] also reveals that whenever concrete formulas are available for the DS currents [6], these do indeed yield expressions for h -tensors which are equivalent to those obtained from our previous formula for k -tensors given above, so that the old and new definitions are entirely consistent. The new approach can be applied to all symmetric space sigma-models too [4].

The commuting quantities in the DS/mKdV hierarchies are naturally identified with the conserved charges of affine Toda field theories (the fields u^i being derivatives of the usual Toda fields). The relationships sketched here underscore the fact that these two prominent categories of integrable Lagrangian field theories, namely sigma-models and Toda theories, have much more in common than might initially be supposed. Although the two classes of models appear to represent radically different ways of introducing interactions amongst collections of fields, we have found that they nevertheless contain a common core, each exhibiting commuting local charges based on precisely the same sets of Weyl-invariant tensors.

Acknowledgements: My warm thanks to the organizers of the conference, Henrik Aratyn and Alexander Sorin, for the invitation to speak. Much of the research described here was carried out while spending a stimulating and very enjoyable year at Princeton University; I am grateful to all in the Physics Department, but especially Erik Verlinde, for making this possible. The work was supported by NSF grant PHY98-02484 and by a PPARC Advanced Fellowship.

References

- [1] J.M. Evans, M. Hassan, N.J. MacKay, and A.J. Mountain, *Local conserved charges in principal chiral models*, Nucl. Phys. **B561** (1999) 385-412; [hep-th/9902008](#).
- [2] J.M. Evans, M. Hassan, N.J. MacKay, and A.J. Mountain, *Conserved charges and supersymmetry in principal chiral and WZW models*, Nucl. Phys. **B580** (2000) 605-646; [hep-th/0001222](#).
- [3] J.M. Evans and A.J. Mountain, *Commuting charges and symmetric spaces*, Phys. Lett. **B483** (2000) 290-298; [hep-th/0003264](#).
- [4] J.M. Evans, *Integrable sigma-models and Drinfeld-Sokolov hierarchies*; [hep-th/0101231](#).
- [5] E. Corrigan, *Recent developments in affine Toda quantum field theory*, Lectures at Banff Summer School 1994; [hep-th/9412213](#).
- [6] V.G. Drinfeld and V.V. Sokolov, *Lie algebras and equations of Korteweg-deVries type*, J. Sov. Math. **30** (1985) 1975-2036.